

Astrometric Measurement of WDS 13433-2458 AB

Roswell Roberts¹, Derek Chow¹, Kevin Lu¹, Marie Yokers², Pat Boyce³, and Grady Boyce³

1. Westview High School, San Diego, California, USA

2. San Diego Mesa College, San Diego, California, USA

3. Boyce Research Initiatives and Education Foundation, San Diego, California, USA

Abstract: By utilizing Las Cumbres Observatory (LCO) telescopes equipped with CCD imaging, measurements of theta and rho of the proposed binary system WDS 13433-2458 AB (HJ 2671) were determined. When compared to the 18 previous measurements of the system however, the data shows no signs of orbital motion and suggests that WDS 13433-2458 AB is not a physical double.

Introduction

By determining the true nature of binary and multiple star systems—whether or not they are gravitationally bound—can assist astronomers in determining the masses of each star and other stellar characteristics such as radius, density, and a mass-luminosity relationship (MLR), can be estimated. As part of the Boyce Research Initiatives and Education Foundation (BRIEF), this research project aims to assist in the verification of whether a system is a gravitationally bound double or an optical double.

WDS 13433-2458 AB (HJ 2671AB) was selected after meeting the following criteria: right ascension (RA) of 12 to 16 hours for reduced air mass during imaging, delta magnitude of 3 or less, and separation of 5 to 15 arcseconds for image clarity.

Materials and Methods

Historical data for HJ 2671 AB was furnished by the United States Naval Observatory, from which we derived a historical chart of theta and rho over time, Figure 1. The first measurement of HJ 2671 was in 1831 and was measured a total of 18 times with the most recent observation being in 2016. Several sources, the Washington Double Star Catalog (WDS), GAIA, and Aladin 10 were consulted for stellar data such as radial velocity, proper motion vectors, and parallax data. Key data points from GAIA are shown in Table 1.

The Las Cumbres Observatory (LCO) 0.4m telescopes were used to observe HJ 2671AB. Requests for these images were processed through the LCO Observ-




Figure 1: Historical Graph

ing Portal with exposure times calculated using the LCO Exposure Time Calculator. SBIG CCD cameras with 2048 X 3072 pixels had a resolution of 0.57" per pixel and a field of view of 19' X 29'. An SDSS-g filter was used to photograph HJ 2671 on Julian Date

Astrometric Measurement of WDS 13433-2458 AB

	Primary Star	Secondary Star
	HJ 2671 A	HJ 2671 B
Proper Motion RA	-33.407	-38.895
Proper Motion RA Error	0.109	0.089
Proper Motion Dec	-14.294	0.8
Proper Motion Dec Error	0.108	0.081
Parallax	7.9587	1.8924
Parallax Error	0.0588	0.047
Radial Velocity	-13.12	-79.31
Radial Velocity Error	0.69	0.35

Table 1: Table of GAIA data

2458580.655185208 for a total of 12 images at 3 seconds exposure time each. With this telescope setting we determined that of 2458580.655185208 HJ 2671 AB had a theta value of 66.099 degrees and a rho value of 27.507 arcseconds.

Each image was processed through the Our Solar Sisters (OSS) Pipeline to remove image imperfections and insert World Coordinate System (WCS) coordinates into the image files (Fitzgerald 2018). AstroImageJ (AIJ) was used to provide measurements of theta and rho (figure 2) in order to compare them to the measurements found in the historical data.

Results

Table 1 gives the data obtained from the GAIA database. Table 2 shows the Mean, Standard Deviation, and Standard Error of the Mean for the position angle (θ) and separation (ρ) measured for HJ 2671AB.

Discussion

A graph of the historical data of the star (figure 1, above) gives a clear pattern in the movement of the B star. The trend indicates that the B star is moving in a Northwesterly direction with a proper motion value of $\langle -38.895, 0.800 \rangle$. However, the A star in the system is not moving with the B star and instead is moving in a Southwesterly direction with a proper motion value of $\langle -33.407, -14.294 \rangle$, which is illustrated in Figure 3 where the proper motion vectors of the A and B star have been superimposed onto the image. Additionally, data from the GAIA satellite reveals that the radial velocities of the A and B star respectively are -13.12 and -79.31. Since both of these values are negative, the two stars are moving away from Earth but because there is a 66.19 difference in radial velocities, we do not believe




Figure 2: Images of HJ 2671 AB being measured in AstroimageJ

Epoch 2019.264	Mean	Std Dev	Std Dev of Mean
θ (deg)	66.099	0.065	0.021
ρ (as)	27.505	0.028	0.009

Table 2. The mean, standard deviation, and standard dev of the mean of the data collected

that they are moving away from Earth together.

Analysis of this data suggests that the stars are not moving together and instead are pursuing different paths through space. This is further supported by parallax data acquired from the GAIA satellite. The A star has a parallax of 7.9587 and an error of 0.0588, which puts the A star at a distance of 406.61 to 412.66 light years away from Earth. The B star's parallax is 1.8924 with a parallax error of 0.0471, placing the B star at a distance of 1681.19 to 1767.03 light years away, making the two stars separated by at least 1268.53 light years. The distance by which these two stars are sepa-

Astrometric Measurement of WDS 13433-2458 AB




Figure 3. An image of HJ 2671 AB with proper motion vectors

rated is considerable and considering the inverse square relationship between the force of gravity and distance, we do not believe that HJ 2671 A is gravitationally bound to HJ 2671 B over a distance of 1268.53 light years.

Conclusion

The large distance between the A and B stars, the lack of a trend indicative of an orbit on the historical graph with our measurements added to it, and the difference in proper motion vectors indicate that the star system is not a physical double. However, inconsistencies with historical data and measurement differences of separation and position angle warrants further observation of the system.

Acknowledgments

We thank Brian Mason and the United States Naval Observatory for providing historical measurement data

and use of the LCO Worldwide telescope systems. We also thank the Boyce Research Initiatives and Education Foundation (BRIEF) for their generous support throughout the study that allowed us to access the remote telescope system and software necessary. Additionally, we thank Michael Fitzgerald at the OSS Pipeline for processing images and removing imperfections made by the camera. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

References

- Collins, K. A., Kielkopf, J. F., Stassun, K. G., and Hessman, F. V., 2018, “AstroImageJ: image processing and photometric extraction for ultra-precise astronomical light curves”, *The Astronomical Journal*, **153**(2), 77.
- Fitzgerald, M.T., 2018, “The Our Solar Siblings Pipeline: Tackling the data issues of the scaling problem for robotic telescope-based astronomy education projects.”, *Robotic Telescopes, Student Research and Education Proceedings*.
- Mason, B. and Hartkopf, W., 2015, *The Washington Double Star Catalog*, Astrometry Department, U.S. Naval Observatory.